

Instructions for Post-EVLT

- 1. Leave pressure bandage on for 24 hours. Do not get the bandage wet.
- 2. After the bandage is removed, you may wash as you normally would.
- 3. For the first three days, apply the antibiotic ointment given to you once to twice daily on the access site located around the knee area.
- 4. You can expect moderate bruising. Discomfort is to be expected. You may take Tylenol Extra-strength for the discomfort.
- 5. Normal daily activity can be resumed. Walking is encouraged throughout the day for the first week after EVLT. Avoid rigorous activities, such as gym workouts.
- 6. You must wear your compression stockings on the treated leg all day and night (except for bathing) for the first week. You will be further instructed at your first post-EVLT visit.
- 7. Please call the office or doctor if you have any concerns or questions.

If you experience any problems or have questions during the healing process, please contact us. Johnson Ferry/Perimeter & Saint Joseph's Hospital – During office hours: 404-257-9933 | After hours: 770-928-5285 Paulding/West Cobb – During office hours: 678-363-3343 | After hours: 770-928-5288

Johnson Ferry/Perimeter Office - 875 Johnson Ferry Rd., NE, Suite 300 - Atlanta, GA 30342 - Phone: 404-257-9933 - Fax: 404-257-9931 Saint Joseph's Hospital Office - 5667 Peachtree Dunwoody Rd., Suite 180 - Atlanta, GA 30342 - Phone: 404-257-9933 - Fax: 404-257-9931 Paulding/ West Cobb Office - 148 Bill Carruth Parkway, Suite 280 - Hiram, Georgia 30141 - Phone: 678-363-3343 - Fax: 678-363-3380